arXiv:1503.05394v1 [math.CA] 14 Feb 2015

A NOTE ON THE NORM CONVERGENCE BY VILENKIN-FEJÉR MEANS

GEORGE TEPHNADZE

Abstract. The main aim of this paper is to find necessary and sufficient conditions for the convergence of Fejér means in terms of the modulus of continuity on the Hardy spaces H_p, when $0 < p \leq 1/2$.

2010 Mathematics Subject Classification. 42C10.

Key words and phrases: Vilenkin system, Fejér means, martingale Hardy space, modulus of continuity.

Let $\mathbb{N}_+$ denote the set of the positive integers, $\mathbb{N} := \mathbb{N}_+ \cup \{0\}$.

Let $m := (m_0,m_1,...)$ denote a sequence of the positive integers not less than 2.

Denote by

$$Z_{m_k} := \{0, 1, ..., m_k - 1\}$$

the additive group of integers modulo m_k.

Define the group G_m as the complete direct product of the group Z_{m_j} with the product of the discrete topologies of Z_{m_j}'s.

The direct product μ of the measures

$$\mu_k(\{j\}) := 1/m_k \qquad (j \in Z_{m_k})$$

is the Haar measure on G_m with $\mu(G_m) = 1$.

If the sequence $m := (m_0,m_1,...)$ is bounded than G_m is called a bounded Vilenkin group, else it is called an unbounded one. **In this paper we discuss bounded Vilenkin groups only.**

The elements of G_m represented by sequences

$$x := (x_0, x_1, ..., x_j, ...) \qquad (x_k \in Z_{m_k}).$$

It is easy to give a base for the neighborhood of G_m

$$I_0(x) := G_m,$$

$$I_n(x) := \{y \in G_m \mid y_0 = x_0, ..., y_{n-1} = x_{n-1}\} \ (x \in G_m, \ n \in \mathbb{N}).$$

Denote $I_n := I_n(0)$, for $n \in \mathbb{N}$ and $\overline{I_n} := G_m \setminus I_n$. Set

$$e_n := (0, ..., 0, 1, 0, ...) \in G_m,$$

the n-th coordinate of which is 1 and the rest are zeros $(n \in \mathbb{N})$.

The research was supported by Shota Rustaveli National Science Foundation grant no.52/54 (Bounded operators on the martingale Hardy spaces).

If we define the so-called generalized number system based on m as $M_0 := 1,\ M_{k+1} := m_k M_k \quad (k \in \mathbb{N})$ then every $n \in \mathbb{N}$ can be uniquely expressed as $n = \sum_{k=0}^{\infty} n_j M_j$, where $n_j \in Z_{m_j} \quad (j \in \mathbb{N})$ and only a finite number of n_j's differ from zero. Let $|n| := \max \{j \in \mathbb{N},\ n_j \neq 0\}$.

Denote by $L_1(G_m)$ the usual (one dimensional) Lebesgue space.

Define the complex valued function $r_k(x) : G_m \to \mathbb{C}$, the generalized Rademacher functions as

$$r_k(x) := \exp(2\pi \iota x_k / m_k) \qquad \left(\iota^2 = -1,\ x \in G_m,\ k \in \mathbb{N}\right)$$

Now define the Vilenkin system $\psi := (\psi_n : n \in \mathbb{N})$ on G_m as:

$$\psi_n(x) := \prod_{k=0}^{\infty} r_k^{n_k}(x) \qquad (n \in \mathbb{N})$$

Specifically, we call this system the Walsh-Paley one if $m \equiv 2$. The Vilenkin system is orthonormal and complete in $L_2(G_m)$ [1, 21].

If $f \in L_1(G_m)$ we can establish the the Fourier coefficients, the partial sums of the Fourier series, the Fejér means, the Dirichlet and Fejér kernels in the usual manner:

$$\begin{aligned}
\widehat{f}(k) &:= \int_G f\overline{\psi_k}\,d\mu, \qquad (\ n \in \mathbb{N}) \\
S_n f &:= \sum_{k=0}^{n-1} \widehat{f}(k)\,\psi_k\ , \quad (n \in \mathbb{N}_+, S_0 f := 0), \\
\sigma_n f &:= \frac{1}{n}\sum_{k=1}^{n} S_k f, \quad D_n := \sum_{k=0}^{n-1} \psi_k\ , \quad K_n := \frac{1}{n}\sum_{k=1}^{n} D_k, \quad (\ n \in \mathbb{N}_+\ ).
\end{aligned}$$

Recall that

$$D_{M_n}(x) = \begin{cases} M_n, & \text{if } x \in I_n \\ 0, & \text{if } x \notin I_n \end{cases} \tag{1}$$

Let $q_A = M_{2A} + M_{2A-2} + \dots + M_2 + M_0,\ 2 < A \in \mathbb{N}_+$. Then (see [2, 9])

$$q_{A-1}\left|K_{q_{A-1}}(x)\right| \geq \frac{M_{2k}M_{2s}}{4}, \text{ for } x \in I_{2s+1}(s_{2k}e_{2k} + s_{2l}e_{2l}), \tag{2}$$

where $1 \leq s_{2k} \leq m_{2k} - 1$ and $1 \leq s_{2s} \leq m_{2s} - 1,\ k = 0, 1, \dots, A-3,$ $s = k+2, k+3, \dots, A-1.$

The norm (or quasinorm) of the space $L_p(G_m)$ is defined by

$$\|f\|_p := \left(\int_{G_m} |f|^p\,d\mu\right)^{1/p} \qquad (0 < p < \infty).$$

The space $L_{p,\infty}(G_m)$ consists of all measurable functions f for which

$$\|f\|_{L_{p,\infty}} := \sup_{\lambda > 0} \lambda^p \mu(f > \lambda) < +\infty$$

The σ-algebra generated by the intervals $\{I_n(x) : x \in G_m\}$ will be denoted by $\mathcal{F}_n\ (n \in \mathbb{N})$. Denote by $f = \left(f^{(n)}, n \in \mathbb{N}\right)$ a martingale with respect to

F_n $(n \in \mathbb{N})$. (for details see e.g. [22]). The maximal function of a martingale f is defend by

$$f^* = \sup_{n\in\mathbb{N}} \left| f^{(n)} \right|.$$

For $0 < p < \infty$ the Hardy martingale spaces H_p (G_m) consist of all martingales, for which

$$\|f\|_{H_p} := \|f^*\|_p < \infty.$$

If $f = \left(f^{(n)},\ n \in \mathbb{N}\right)$ is martingale then the Vilenkin-Fourier coefficients must be defined in a slightly different manner:

$$\widehat{f}\,(i) := \lim_{k\to\infty} \int_{G_m} f^{(k)}\,(x)\,\overline{\Psi}_i\,(x)\,d\mu\,(x)\,.$$

The concept of modulus of continuity in H_p $(0 < p \leq 1)$ is defined by

$$\omega\,(1/M_n, f)_{H_p} := \|f - S_{M_n} f\|_{H_p}\,.$$

A bounded measurable function a is p-atom, if there exist a dyadic interval I, such that

$$\int_I a d\mu = 0, \quad \|a\|_\infty \leq \mu\,(I)^{-1/p}\,, \quad \operatorname{supp}(a) \subset I.$$

The dyadic Hardy martingale spaces H_p (G) for $0 < p \leq 1$ have an atomic characterization. Namely the following theorem is true (see [24]):

Theorem W. A martingale $f = (f_n,\ n \in \mathbb{N})$ is in $H_p\,(0 < p \leq 1)$ if and only if there exists a sequence $(a_k,\ k \in \mathbb{N})$ of p-atoms and a sequence $(\mu_k,\ k \in \mathbb{N})$ of a real numbers such that for every $n \in \mathbb{N}$

$$\sum_{k=0}^{\infty} \mu_k S_{M_n} a_k = f_n \tag{3}$$

and

$$\sum_{k=0}^{\infty} |\mu_k|^p < \infty,$$

Moreover, $\|f\|_{H_p} \backsim \inf \left(\sum_{k=0}^{\infty} |\mu_k|^p\right)^{1/p}$, where the infimum is taken over all decomposition of f of the form (3).

The weak type-$(1,1)$ inequality for maximal operator of Fejér means σ^* can be found in Schipp [14] for Walsh series and in Pál, Simon [13] for bounded Vilenkin series. Fujji [5] and Simon [16] verified that σ^* is bounded from H_1 to L_1. Weisz [23] generalized this result and proved the boundedness of σ^* from the martingale space H_p to the space L_p for $p > 1/2$. Simon [15] gave a counterexample, which shows that boundedness does not hold for $0 < p < 1/2$. The counterexample for $p = 1/2$ due to Goginava [6], (see also [3] and [10]). Weisz [25] proved that σ^* is bounded from the Hardy space $H_{1/2}$ to the space $L_{1/2,\infty}$.

The author in [18] and [19] (for Walsh system see [7]) proved that the maximal operator $\widetilde{\sigma}_p^*$ with respect Vilenkin system defined by

$$\widetilde{\sigma}_p^* := \sup_{n\in\mathbb{N}} \frac{|\sigma_n|}{(n+1)^{1/p-2}\log^{2[1/2+p]}(n+1)},$$

where $0<p\leq 1/2$ and $[1/2+p]$ denotes integer part of $1/2+p$, is bounded from the Hardy space H_p to the space L_p. Moreover, we showed that the order of deviant behaviour of the n-th Fejér means was given exactly. As a corollary we get

$$\|\sigma_n f\|_p \leq c_p\,(n+1)^{1/p-2}\log^{2[1/2+p]}(n+1)\,\|f\|_{H_p}. \tag{4}$$

For Walsh-Kaczmarz system analogical theorems are proved in [8] and [20].

Móricz and Siddiqi [12] investigates the approximation properties of some special Nörlund means of Walsh-Fourier series of L_p function in norm. Fridly, Manchanda and Siddiqi [4] improve and extend the results of Móricz and Siddiqi [12] among them in H_p norm, where $0<p<1$. In [11] Goginava investigated the behavior of Cesáro means of Walsh-Fourier series in detail.

The main aim of this paper is to find necessary and sufficient conditions for the convergence of Fejér means in terms of the modulus of continuity on the Hardy spaces H_p, when $0<p\leq 1/2$. In particular, the following is true:

Theorem 1. *Let* $0<p\leq 1/2,\ f\in H_p,\ M_N<n\leq M_{N+1}$ *and*

$$\omega\left(\frac{1}{M_N},f\right)_{H_p} = o\left(\frac{1}{M_N^{1/p-2}N^{2[1/2+p]}}\right),\ \ \text{as}\ \ N\to\infty. \tag{5}$$

Then

$$\|\sigma_n f - f\|_p \to 0,\ \ \text{when}\ n\to\infty.$$

Theorem 2. *a) Let* $0<p<1/2$ *and* $M_N<n\leq M_{N+1}$. *There exists a martingale* $f\in H_p(G_m)$ *for which*

$$\omega\left(\frac{1}{M_N},f\right)_{H_p} = O\left(\frac{1}{M_N^{1/p-2}}\right),\quad \text{as}\ \ N\to\infty \tag{6}$$

and

$$\|\sigma_n f - f\|_{L_{p,\infty}} \nrightarrow 0,\ \ \text{as}\ \ n\to\infty.$$

b) Let $M_N<n\leq M_{N+1}$. *There exists a martingale* $f\in H_{1/2}(G_m)$ *for which*

$$\omega\left(\frac{1}{M_N},f\right)_{H_{1/2}} = O\left(\frac{1}{N^2}\right),\quad \text{as}\ \ N\to\infty \tag{7}$$

and

$$\|\sigma_n f - f\|_{1/2} \nrightarrow 0,\ \ \text{as}\ \ n\to\infty.$$

Proof of Theorem 1. Let $f \in H_p$, $0 < p \leq 1/2$ and $M_N < n \leq M_{N+1}$. Using (4) we have

$$\begin{aligned}
& \|\sigma_n f - f\|_p^p \\
\leq\ & \|\sigma_n f - \sigma_n S_{M_N} f\|_p^p + \|\sigma_n S_{M_N} f - S_{M_N} f\|_p^p + \|S_{M_N} f - f\|_p^p \\
=\ & \|\sigma_n (S_{M_N} f - f)\|_p^p + \|S_{M_N} f - f\|_p^p + \|\sigma_n S_{M_N} f - S_{M_N} f\|_p^p \\
\leq\ & c_p \left(n^{1-2p} \log^{2p[1/2+p]} n + 1\right) \omega^p \left(\frac{1}{M_N}, f\right)_{H_p} + \|\sigma_n S_{M_N} f - S_{M_N} f\|_p^p .
\end{aligned}$$

By simple calculation we get

$$\sigma_n S_{M_N} f - S_{M_N} f = \frac{M_N}{n} \left(\sigma_{M_N} S_{M_N} f - S_{M_N} f\right).$$

Let $p > 0$ and $f \in H_p$. Since (see [22] (for bounded Vilenkin systems))

$$\|\sigma_{M_k} f - f\|_p \to 0, \quad \text{when } k \to \infty \tag{8}$$

$S_{M_N} \sigma_{M_N} f = \sigma_{M_N} S_{M_N} f$ and $S_{M_N} f$ is martingale, we obtain

$$\|\sigma_n S_{M_N} f - S_{M_N} f\|_p^p = \frac{M_N^p}{n^p} \|\sigma_{M_N} S_{M_N} f - S_{M_N} f\|_p^p \to 0, \text{ when } k \to \infty. \tag{9}$$

It follows that under condition (5) we have

$$\|\sigma_n f - f\|_p \to 0, \text{ when } n \to \infty.$$

Which complete the proof of Theorem 1.

Proof of Theorem 2. At first we consider case $0 < p < 1/2$. Let

$$a_k = \frac{M_k^{1/p-1}}{\lambda} \left(D_{M_k+1} - D_{M_k}\right),$$

where $\lambda = \sup_{n \in \mathbb{N}} m_n$. It is easy to show that a_k are p-atoms.

We set

$$f^{(A)} = \sum_{i=0}^{A} \frac{\lambda}{M_i^{1/p-2}} a_i.$$

Using Theorem W (see [17]) we conclude that martingale $f \in H_p$.

On the other hand

$$\begin{aligned}
& f - S_{M_n} f \\
=\ & \left(f^{(1)} - S_{M_n} f^{(1)}, ..., f^{(n)} - S_{M_n} f^{(n)}, ..., f^{(n+k)} - S_{M_n} f^{(n+k)}\right) \\
=\ & \left(0, ..., 0, f^{(n+1)} - f^{(n)}, ..., f^{(n+k)} - f^{(n)}, ...\right) \\
=\ & \left(0, ..., 0, \sum_{i=n+1}^{n+k} \frac{a_i}{M_i^{1/p-2}}, ...\right), \quad k \in \mathbb{N}_+
\end{aligned} \tag{10}$$

is martingale. Applying Theorem W and (10) we get

$$\omega(\frac{1}{M_n},f)_{H_p}\leq\sum_{i=n+1}^{\infty}\frac{1}{M_i^{1/p-2}}\leq\frac{c}{M_n^{1/p-2}}=O\left(\frac{1}{M_n^{1/p-2}}\right).$$

By simple calculation we have

$$\widehat{f}(j)=\begin{cases}M_i, & \text{if } j\in\{M_i,...,M_{i+1}-1\}\,,\ i=0,1,...\\ 0, & \text{if } j\notin\bigcup\limits_{i=0}^{\infty}\{M_i,...,M_{i+1}-1\}\,.\end{cases}\tag{11}$$

Combining (8) and (11) we can write

$$\begin{aligned}&\limsup_{k\to\infty}\|\sigma_{M_k+1}f-f\|_{L_{p,\infty}}\\ =\ &\limsup_{k\to\infty}\|\frac{M_k\sigma_{M_k}f}{M_k+1}+\frac{S_{M_k}(f)}{M_k+1}+\frac{M_kw_{M_k+1}}{M_k+1}-\frac{M_kf}{M_k+1}-\frac{f}{M_k+1}\|_{L_{p,\infty}}\\ \geq\ &\limsup_{k\to\infty}\frac{M_k}{M_k+1}\|w_{M_k}\|_{L_{p\infty}}\\ &-\limsup_{k\to\infty}\frac{M_k}{M_k+1}\|\sigma_{M_k}f-f\|_{L_{p,\infty}}-\limsup_{k\to\infty}\frac{1}{M_k+1}\|S_{M_k}f-f\|_{L_{p,\infty}}\\ \geq\ &\limsup_{k\to\infty}\frac{M_k}{M_k+1}\geq c>0.\end{aligned}$$

Now prove second part of Theorem 2. Let

$$f^{(A)}=\sum_{i=1}^{A}\frac{\lambda}{M_i^2}a_i,$$

where

$$a_i=\frac{M_{2M_i}}{\lambda}\left(D_{M_{2M_i}+1}-D_{M_{2M_i}}\right)$$

and $\lambda=\sup_{n\in\mathbb{N}}m_n$.

Analogously, we conclude that $f\in H_{1/2}$ and

$$\omega(\frac{1}{M_n},f)_{H_{1/2}}\leq\sum_{i=[\log n/2]}^{A}\frac{1}{M_i^2}=O\left(\frac{1}{n^2}\right).$$

Hence

$$\sigma_{q_{M_k}}f-f=\frac{M_{2M_k}\sigma_{M_{2M_k}}(f)}{q_{M_k}}+\frac{1}{q_{M_k}}\sum_{j=M_{2M_k}+1}^{q_{M_k}}S_jf-\frac{M_{2M_k}f}{q_{M_k}}-\frac{q_{M_k-1}f}{q_{M_k}}\tag{12}$$

It is easy to show that

$$\widehat{f}(j)=\begin{cases}\frac{M_{2M_i}}{M_i^2}, & \text{if } j\in\{M_{2M_i},...,M_{2M_i+1}-1\}\,,\ i=0,1,...\\ 0, & \text{if } j\notin\bigcup\limits_{i=0}^{\infty}\{M_{2M_i},...,M_{2M_i+1}-1\}\,.\end{cases}\tag{13}$$

Let $M_{2M_k} < j \leq q_{M_k}$. Since $D_{j+M_{2M_k}} = D_{M_{2M_k}} + w_{M_{2M_k}} D_j$, when $j < M_{2M_k}$ using (13) we have

$$S_j f = S_{M_{2M_k}} f + \sum_{v=M_{2M_k}}^{j-1} \widehat{f}(v) w_v = S_{M_{2M_k}} f + \frac{M_{2M_k} w_{M_{2M_k}} D_{j-M_{2M_k}}}{M_k^2}.$$

and

$$\frac{1}{q_{M_k}} \sum_{j=M_{2M_k}+1}^{q_{M_k}} S_j f = \frac{q_{M_k-1} S_{M_{M_k}} f}{q_{M_k}} + \frac{M_{2M_k} w_{M_{2M_k}}}{q_{M_k} M_k^2} \sum_{j=1}^{q_{M_k-1}} D_j$$

$$= \frac{q_{M_k-1} S_{M_{2M_k}} f}{q_{M_k}} + \frac{M_{2M_k} w_{M_{2M_k}} q_{M_k-1} K_{q_{M_k-1}}}{q_{M_k} M_k^2}.$$

Using (12) we get

$$\|\sigma_{q_{M_k}} f - f\|_{1/2}^{1/2} \geq \frac{c}{M_k} \|q_{M_k-1} K_{q_{M_k-1}}\|_{1/2}^{1/2} \tag{14}$$

$$- \left(\frac{M_{2M_k}}{q_{M_k}}\right)^{1/2} \|\sigma_{M_{2M_k}} f - f\|_{1/2}^{1/2} - \left(\frac{q_{M_k-1}}{q_{M_k}}\right)^{1/2} \|S_{M_{2M_k}} f - f\|_{1/2}^{1/2}.$$

Let $x \in I_{2s+1}\left(s_{2k} e_{2k} + s_{2\eta} e_{2\eta}\right), 1 \leq s_{2\eta} \leq m_{2\eta} - 1$ and $1 \leq s_{2k} \leq m_{2k} - 1$, $\eta = 1, ..., M_k - 4,\ s = \eta + 2, \eta + 3, ..., M_k - 2$. Applying (2) we have

$$q_{M_k-1} \left| K_{q_{M_k-1}}(x) \right| \geq \frac{M_{2\eta} M_{2s}}{4}.$$

We can write

$$\int_{G_m} \left| q_{M_k-1} K_{q_{M_k-1}} \right|^{1/2} d\mu \tag{15}$$

$$\geq \frac{1}{4} \sum_{\eta=1}^{M_k-4} \sum_{s=\eta+2}^{M_k-2} \int_{I_{2s+1}(s_{2k}e_{2k}+s_{2l}e_{2l})} \left| q_{M_k-1} K_{q_{M_k-1}} \right|^{1/2} d\mu$$

$$\geq \frac{1}{4} \sum_{\eta=1}^{M_k-4} \sum_{s=\eta+2}^{M_k-2} \frac{1}{2M_{2s}} \sqrt{M_{2s} M_{2\eta}} \geq c M_k.$$

Combining (8), (14-15) we have

$$\limsup_{k\to\infty} \|\sigma_{q_{M_k}} f - f\|_{1/2} \geq c > 0.$$

Theorem 2 is proved.

References

[1] Agaev, G. N., Vilenkin, N. Ya., DzhsafarlyG. M., Rubinshtein, A. I.: Multiplicative systems of functions and harmonic analysis on zero-dimensional groups, Baku, Ehim, 1981 (in Russian).

[2] Blahota, I., Gát, G., G., Goginava, U.: maximal operators of Fejer means of double Vilenkin-Fourier series, Colloq. Math. 107 (2007), no. 2, 287–296.

[3] Blahota, I., Gát, G., G., Goginava, U.: Maximal operators of Fejér means of Vilenkin-Fourier series. JIPAM. J. Inequal. Pure Appl. Math. 7 (2006), 1-7.

[4] Fridly, S., Manchanda, P., Siddiqi, A.: Approximation by Walsh-Nörlund means, Acta Sci. Math. (Szeged) 74, 3-4, (2008), 593-608.

[5] Fujii, N. J.: A maximal inequality for H_1 functions on the generalized Walsh-Paley group, Proc. Amer. Math. Soc. 77 (1979), lll-116.

[6] Goginava, U.: The maximal operator of Marcinkiewicz-Fejér means of the d-dimensional Walsh-Fourier series. East J. Approx. 12 (2006), no. 3, 295–302.

[7] Goginava, U.: Maximal operators of Fejér-Walsh means. Acta Sci. Math. (Szeged) 74 (2008), no. 3-4, 615–624.

[8] Goginava, U., NAGY, K.: On the maximal operator of Walsh-Kaczmarz-Fejer means, Czechoslovak Math. J. 61(136), 2011, 673-686.

[9] Goginava, U.: Maximal operators of Fejér means of double Walsh-Fourier series. Acta Math. Hungar. 115 (2007), no. 4, 333–340.

[10] Goginava, U.: The maximal operator of the Fejér means of the character system of the p-series field in the Kaczmarz rearrangement. Publ. Math. Debrecen 71 (2007), no. 1-2, 43–55.

[11] Goginava, U.: On the approximation properties of Cesáro means of negative order of Walsh-Fourier series. Journal of approximation theory 115, 9-20 (2002).

[12] Möricz, F., Siddiqi, A.: Approximation by Nörlund means of Walsh-Fourier series. Journal of approximation theory. 70 1992, (375-389).

[13] Pál J., Simon, P.: On a generalization of the comncept of derivate, Acta Math. Hung., 29 (1977), 155-164.

[14] Schipp, F.: Certain rearranngements of series in the Walsh series, Mat. Zametki, 18 (1975), 193-201.

[15] Simon, P.: Cesáro summability with respect to two-parameter Walsh sistems, Monatsh. Math .,131 (2000),321-334.

[16] Simon, P.: Inverstigations wish respect to the Vilenkin sistem, Annales Univ. Sci. Budapest Eotv., Sect. Math., 28 (1985) 87-101.

[17] Tephnadze, G.: Fejér means of Vilenkin-Fourier series, Stud. sci. math. Hung., 49 (1), (2012) 79-90.

[18] Tephnadze, G.: On the maximal operator of Vilenkin-Fejér means, Turk. J. Math, (to appear).

[19] Tephnadze, G.: On the maximal operators of Vilenkin-Fejér means on Hardy spaces, Mathematical Inequalities & Applications, 16, no. 2 (2013), 301-312.

[20] Tephnadze, G.: On the maximal operators of Walsh-Kaczmarz-Fejer means, Periodica Math. Hung. (to appear).

[21] Vilenkin, N. Ya.: A class of complate ortonormal systems, Izv. Akad. Nauk. U.S.S.R., Ser. Mat., 11 (1947), 363-400.

[22] Weisz, F.: Martingale Hardy spaces and their applications in Fourier Analysis, Springer, Berlin-Heideiberg-New York, 1994.

[23] Weisz, F.: Cesáro summability of one and two-dimensional Fourier series, Anal. Math. Studies, 5 (1996), 353-367.

[24] Weisz, F.: Hardy spaces and Cesáro means of two-dimensional Fourier series, Bolyai Soc. math. Studies, (1996), 353-367.

[25] Weisz, F.: Weak type inequalities for the Walsh and bounded Ciesielski systems. Anal. Math. 30 (2004), no. 2, 147–160.

G. Tephnadze, Department of Mathematics, Faculty of Exact and Natural Sciences, Iv. Javaxikhishvili, Tbilisi State University, Chavchavadze str. 1, Tbilisi 0128, Georgia

E-mail address: `giorgitephnadze@gmail.com`